

HYPOMONOTONICITY OF THE NORMAL CONE AND PROXIMAL SMOOTHNESS

G.M. IVANOV

ABSTRACT. In this paper we study the properties of the normal cone to the proximally smooth set. We give the complete characterization of a proximally smooth set through the monotony properties of its normal cone in an arbitrary uniformly convex and uniformly smooth Banach space. We give the exact bounds for right-hand side in the monotonicity inequality for normal cone in terms of the moduli of smoothness and convexity of a Banach space.

1. INTRODUCTION

Let X be a real Banach space. For a set $A \subset X$ by ∂A , $\text{int } A$ we denote the boundary and the interior of A . We use $\langle p, x \rangle$ to denote the value of functional $p \in X^*$ at vector $x \in X$. For $R > 0$ and $c \in X$ we denote by $\mathfrak{B}_R(c)$ the closed ball with center c and radius R , by $\mathfrak{B}_R^*(c)$ we denote the ball in the conjugate space. The distance from a point $x \in X$ to a set $A \subset X$ is defined as

$$\rho(x, A) = \inf_{a \in A} \|x - a\|.$$

The *metric projection* of a point x onto a set A is defined as any element of the set

$$P_A(x) = \{a \in A : \|a - x\| = \rho(x, A)\}.$$

Clarke, Stern and Wolenski [1] introduced and studied the *proximally smooth* sets in a Hilbert space H .

Definition 1. A set $A \subset X$ is said to be *proximally smooth* with constant R if the distance function $x \rightarrow \rho(x, A)$ is continuously differentiable on set

$$U(R, A) = \{x \in X : 0 < \rho(x, A) < R\}.$$

We denote by $\Omega_{PS}(R)$ the set of all closed proximally smooth sets with constant R in X .

Properties of proximally smooth sets in a Banach space and the relation between such sets and akin classes of sets were investigated in [1]-[9]. Such sets are usually called weakly convex sets. Some notes about the history of this problem one could find in paper [3]. Perhaps, the most complete description of the proximally smooth sets and their properties in a Hilbert space can be found in [5].

Definition 2. The normal cone to a set $A \subset X$ at a point $a_0 \in A$ is defined as follows

$$N(a_0, A) = \left\{ p \in X^* : \forall \varepsilon > 0 \exists \delta > 0 : \quad \forall a \in A \cap \mathfrak{B}_\delta(a_0) \quad \langle p, a - a_0 \rangle \leq \varepsilon \|a - a_0\| \right\}.$$

Remark 1. It is easily seen that normal cone $N(a_0, A)$ does not change when the norm of the space X changes in an equivalent way.

The problem of characterization of proximally smooth sets in Hilbert spaces through the monotony properties of the normal cone turns out to be very important for applications. Thus, Theorem 1.9.1 of monograph [5] may be reformulated in the following way (See also [3], Cor. 2.2).

Theorem I. *Let A be a closed set in a Hilbert space H and $R > 0$. The following conditions are equivalent*

- (1) *the set A is proximally smooth with constant $R > 0$;*
- (2) *for any vectors $x_1, x_2 \in A$, $p_1 \in N(x_1, A)$, $p_2 \in N(x_2, A)$ such that $\|p_1\| = \|p_2\| = 1$, the following inequality holds*

$$(1) \quad \langle p_2 - p_1, x_2 - x_1 \rangle \geq -\frac{\|x_2 - x_1\|^2}{R}.$$

The equivalent in a Hilbert space properties of the normal cone were studied earlier in [1], [2]. Note that the fact that inequality (1) holds for proximal smooth sets in Hilbert spaces was the starting point for the construction of the proximal subgradient, which is an important construction in optimization nowadays.

Afterwards, of course, attempts were made to extend this result to Banach spaces. Here we must note the works of Thibault and his colleagues, for example, [6], [7]. As a rule, in these works other approaches to defining the normal cone are examined, and then the coincidence of different normal cones for proximally smooth sets in special Banach spaces (usually, uniformly smooth and uniformly convex) is proved.

Perhaps, the strongest results in this domain may be formulated in the following two theorems.

Theorem II. *Let X be a uniformly convex and uniformly smooth Banach space. Let $\rho_X(\tau) \asymp \tau^2$ as $\tau \rightarrow 0$. Then a proximally smooth set $A \subset X$ with constant $r > 0$ satisfies condition 2) of Theorem I for some constant $R > 0$.*

Theorem III. *Let the convexity and smoothness moduli be of power order at zero in a Banach space X . Let $\delta_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$. Then, if a set A satisfies condition 2) of Theorem I, then it is proximally smooth with some constant $r > 0$.*

In this paper, using the geometrical properties of the sets from the class $\Omega_{PS}(R)$, the connection between the smoothness of a set and the properties of its normal cone is studied.

2. MAIN RESULTS

In the sequel we shall need some additional notation.

By definition, put $J_1(x) = \{p \in \partial \mathfrak{B}_1^*(o) : \langle p, x \rangle = \|x\|\}$.

Let

$$\delta_X(\varepsilon) = \inf \left\{ 1 - \frac{\|x + y\|}{2} : x, y \in \mathfrak{B}_1(0), \|x - y\| \geq \varepsilon \right\}$$

and

$$\rho_X(\tau) = \sup \left\{ \frac{\|x + y\|}{2} + \frac{\|x - y\|}{2} - 1 : \|x\| = 1, \|y\| = \tau \right\}.$$

The functions $\delta_X(\cdot) : [0, 2] \rightarrow [0, 1]$ and $\rho_X(\cdot) : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are referred to as the moduli of convexity and smoothness of X respectively.

Normed space X is called *uniformly convex*, if $\delta_X(\varepsilon) > 0$ for all $\varepsilon \in (0, 2]$. Normed space X is called *uniformly smooth*, if $\lim_{\tau \rightarrow +0} \frac{\rho_X(\tau)}{\tau} = 0$.

Let f and g be two non-negative functions, each one defined on a segment $[0, \varepsilon]$. We shall consider f and g as *equivalent at zero*, denoted by $f(t) \asymp g(t)$ as $t \rightarrow 0$, if there exist positive constants a, b, c, d, e such that $af(bt) \leq g(t) \leq cf(dt)$ for $t \in [0, e]$.

First we generalize condition 2) of Theorem I.

Definition 3. Let a function $\psi : [0, +\infty) \rightarrow [0, +\infty)$ be given. A set $A \subset X$ satisfies the ψ -hypomonotonicity condition of the normal cone with constant $R > 0$ if for some $\varepsilon > 0$ and for any $x_1, x_2 \in A$, $p_1 \in N(x_1, A)$, $p_2 \in N(x_2, A)$, $\|p_1\| = \|p_2\| = 1$ such that $\|x_1 - x_2\| \leq \varepsilon$, the inequality

$$(2) \quad \langle p_2 - p_1, x_2 - x_1 \rangle \geq -R\psi\left(\frac{\|x_2 - x_1\|}{R}\right)$$

holds. Through $\Omega_N^\psi(R)$ we denote the class of all closed sets $A \subset X$ that satisfy the ψ -hypomonotonicity condition with constant $R > 0$.

Naturally, studying the classes of sets that satisfy the ψ -hypomonotonicity condition, we suppose some restrictions on the function ψ .

Definition 4. Through $\mathfrak{M}$ denote the class of convex and Lipschitz functions $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\psi(0) = 0$.

We say that the function $N : [0, +\infty) \rightarrow [0, +\infty)$ such that $N(0) = 0$, satisfies the *Figiel condition* if there exists a constant K such that the function $N(\cdot)$ on some interval $(0, \varepsilon)$ satisfies the condition

$$(3) \quad \frac{N(s)}{s^2} \leq K \frac{N(t)}{t^2} \quad \forall 0 < t \leq s < \varepsilon.$$

Remark 2. The modulus of smoothness of an arbitrary Banach space satisfies the Figiel condition (see[10], Proposition 1.e.5.).

Definition 5. Through $\mathfrak{M}_2$ denote the class of functions from $\mathfrak{M}$ that satisfy the Figiel condition.

It is easy to see that an arbitrary function $f(\cdot) \in \mathfrak{M}_2$ that is not identically zero satisfies the condition $t^2 = O(f(t))$ as $t \rightarrow 0$.

Obviously, in case of a Hilbert space H for the function $\psi(t) = t^2$ the assertion of Theorem I is equivalent to the equality $\Omega_N^\psi(R) = \Omega_{PS}(R)$.

In the author's opinion, the condition $\psi \in \mathfrak{M}_2$ is rather natural. Moreover, it would be interesting to prove or contradict the following hypothesis.

Conjecture 1. If in a uniformly convex and uniformly smooth Banach space X a set $A \subset X$ belongs to the class $\Omega_{PS}(R)$ and to the class $\Omega_N^\psi(r)$ for some function $\psi \in \mathfrak{M} \setminus \mathfrak{M}_2$ and constants $R > 0, r > 0$, then it belongs to the class $\Omega_N^{\psi_1}(r)$, where $\psi_1(t) = ct^2$ for some $c \geq 0$.

The following two theorems provide an ample description of the connection of classes $\Omega_N^\psi(R), \Omega_{PS}(R)$.

Theorem 1. *In a uniformly convex and uniformly smooth Banach space X the following statements are equivalent for the function $\psi \in \mathfrak{M}$:*

- (1) *there exists $k_1 > 0$ such that $\Omega_{PS}(R) \subset \Omega_N^{k_1\psi}(R)$ for any $R > 0$;*
- (2) *$\rho_X(\tau) = O(\psi(\tau))$ as $\tau \rightarrow 0$.*

Theorem 2. *In a uniformly convex and uniformly smooth Banach space X the following statements are equivalent for the function $\psi \in \mathfrak{M}$:*

- (1) *there exists $k_2 > 0$ such that $\Omega_N^{k_2\psi}(R) \subset \Omega_{PS}(R)$ for any $R > 0$;*
- (2) *$\psi(\varepsilon) = O(\delta_X(\varepsilon))$ as $\varepsilon \rightarrow 0$.*

Clearly, $A \in \Omega_{PS}(1)$ iff $RA \in \Omega_{PS}(R)$ for some $R > 0$. A similar property holds for the class of ψ -hypomonotonous sets. Therefore, Theorems 1, 2 can be proved assuming that R is a fixed positive number. Using Theorems 2, 1 we will show that the following statement, which gives a complete answer to the question about the possibility of the inclusion $\Omega_N^{k_1\psi}(R) \subset \Omega_{PS}(R) \subset \Omega_N^{k_2\psi}(R)$, holds.

Theorem 3. *If in a uniformly smooth and uniformly convex Banach space X for some function $\psi \in \mathfrak{M}$ there exist $k_1 > 0$, $k_2 > 0$ such that the inclusions $\Omega_N^{k_1\psi}(R) \subset \Omega_{PS}(R) \subset \Omega_N^{k_2\psi}(R)$ hold, then $\delta_X(\varepsilon) \asymp \rho_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$, and, therefore, the space X is isomorphic to a Hilbert space.*

Conjecture 2. *The equality $\Omega_{PS}(R) = \Omega_N^\psi(R)$ holds only in a Hilbert space provided that $\psi(t) = t^2$.*

3. PRELIMINARIES

Geometrical properties of class $\Omega_{PS}(R)$ hide in its definition. To clarify the geometrical properties of $\Omega_{PS}(R)$, which are very useful in this paper, we introduce two equivalent (in certain spaces!) definitions of class $\Omega_{PS}(R)$.

Definition 6. *We say that a closed set A in a Banach space X satisfies the P -supporting condition of weak convexity with constant $R > 0$ if the facts that $u \in U(R, A)$ and $x \in P_A(u)$ imply the inequality*

$$\rho\left(x + \frac{R}{\|u - x\|}(u - x), A\right) \geq R.$$

The set of all such sets we denote by $\Omega_P(R)$.

Definition 7. *We say that a closed set A in a Banach space X satisfies the N -supporting condition of weak convexity with constant $R > 0$ if the facts that $x \in A$, $p \in N(x, A) \cap \partial\mathfrak{B}_1^*(o)$ and $u \in \partial\mathfrak{B}_1(o)$ such that $p \in J_1(u)$ imply the inequality $\rho(x + Ru, A) \geq R$. The set of all such sets we denote by $\Omega_N(R)$.*

The equality $\Omega_{PS}(R) = \Omega_P(R)$ was proved in [9] in case of uniformly convex and uniformly smooth Banach space. The equality $\Omega_P(R) = \Omega_N(R)$ was proved in [8] in case of uniformly convex Banach space. So we can formulate the following theorem.

Theorem IV. *Let X be a uniformly smooth and uniformly convex Banach space. Then*

$$\Omega_{PS}(R) = \Omega_P(R) = \Omega_N(R).$$

Further we will use class $\Omega_N(R)$ instead of $\Omega_{PS}(R)$, as mentioned above these classes coincide under conditions of theorems 1, 2, 3.

Theorem 4. *Let X be a reflexive Banach space, $R > 0$, $A \in \Omega_N(R)$, $a_0 \in A$, $p \in N(a_0, A) \cap \partial\mathfrak{B}_1^*(o)$, $\delta > 0$, $\varepsilon = \frac{2R}{\delta} \rho_X\left(\frac{\delta}{R}\right)$. Then*

$$\langle p, a - a_0 \rangle \leq \varepsilon \|a - a_0\| \quad \forall a \in A \cap \mathfrak{B}_\delta(a_0).$$

Proof.

Fix a vector $a \in A \cap \mathfrak{B}_\delta(a_0)$. By reflexivity of X , there exists a vector $u \in \partial\mathfrak{B}_1(o)$ such that $p \in J_1(u)$. Since $A \in \Omega_N(R)$, we have that $\rho(a_0 + Ru, A) \geq R$. Therefore, $\|a_0 + Ru - a\| \geq R$. Denote $w = \frac{a - a_0}{R}$. Then $\|u - w\| \geq 1$. By the definition of the modulus of smoothness we get that $2\rho_X(\|w\|) \geq \|u + w\| + \|u - w\| - 2$. Consequently,

$$2\rho_X(\|w\|) \geq \|u + w\| - 1 \geq \langle p, u + w \rangle - 1 = \langle p, w \rangle.$$

And now we have

$$\langle p, a - a_0 \rangle = R\langle p, w \rangle \leq 2R\rho_X(\|w\|) = 2R\rho_X\left(\frac{\|a - a_0\|}{R}\right).$$

Since function $\tau \rightarrow \frac{\rho_X(\tau)}{\tau}$ is increasing and $\|a - a_0\| \leq \delta$, we have $\rho_X\left(\frac{\|a - a_0\|}{R}\right) \leq \frac{\|a - a_0\|}{\delta} \rho_X\left(\frac{\delta}{R}\right)$. Consequently, $\langle p, a - a_0 \rangle \leq \varepsilon \|a - a_0\|$. $\square$

We need to formulate some properties of the moduli of smoothness and convexity.

It is well known ([18] Ch.3, §4, Lemma 1.), that in a Banach space the following inequalities hold.

$$(4) \quad 2 \leq \limsup_{\tau \rightarrow 0} \frac{\rho_X(2\tau)}{\rho_X(\tau)} \leq 4.$$

The following two lemmas are technical and obvious. The first one can find in [19].

Lemma 1. *Let $x, y \in X$, $x \neq 0$, $p \in J_1(x)$. Then*

$$(5) \quad \|x + y\| \leq \|x\| + \langle p, y \rangle + 2\|x\| \rho_X\left(\frac{\|y\|}{\|x\|}\right).$$

Lemma 2. *Let X be a uniformly convex Banach space, $x_0 \in \partial\mathfrak{B}_R(o)$ and $p_0 \in \partial\mathfrak{B}_1^*(o)$ be a functional dual to the vector $-x_0$. Then for any vector $z \in \text{int } \mathfrak{B}_R(o)$ we have*

$$(6) \quad \langle p_0, z - x_0 \rangle > 2R\delta_X\left(\frac{\|z - x_0\|}{R}\right).$$

The proof of the next proposition one could find in work [8].

Theorem V. *Let $A \subset X$, $x_1 \in X \setminus A$, $x_0 \in P_A(x_1)$, and the norm is Frechet differentiable at point $x_1 - x_0$. Then $J_1(x_1 - x_0) \subset N(x_0, A)$.*

We will denote by $\mathfrak{B}_1^p(o)$ the intersection of the unit ball and the hyperplane

$$H_p = \{x \in X \mid \langle p, x \rangle = 0\},$$

i.e. the set $\mathfrak{B}_1(o) \cap H_p$. Let $\mathfrak{B}_1^p(a) = \mathfrak{B}_1^p(o) + oa$.

Lemma 3. *Let $x \in \partial\mathfrak{B}_1(o)$, $p \in J_1(ox)$, $z \in \mathfrak{B}_1^p(x)$. Let a point y such that segment yz is parallel to ox and intersects the unit sphere at unique point y . Then $2\|zx\| \geq \|xy\|$.*

Proof.

By the triangle inequality, it suffices to show that $\|zx\| \geq \|zy\|$. By d denote a point on unit sphere such that vector od is collinear to the vector xz . Let line l be parallel to ox and $d \in l$. By constructions, we have that points x, y, z, o, d and line l lies at the same plane – linear span of the vectors ox and xz . So lines l and xz intersect, by c we denote their intersection point. Note that $odcx$ is a parallelogram and $\|dc\| = 1$; segment dx belongs to the unit ball and does not intersect the interior of the segment zy . Let $y' = zy \cap dx$. By similarity, we have

$$\|zy\| \leq \|zy'\| = \frac{\|xz\|}{\|xc\|} \|dc\| = \|xz\|.$$

□

It is worth noticing, that in the conditions of Lemma 3 we have that z is a projection along vector ox of the point y on some supporting hyperplane to the unit ball at x . Moreover, z belongs to the metrics projection of the point y on this hyperplane. That is to say, Lemma 3 show us that if one projects along vector ox segment xy on the hyperplane, which is supporting to the unit ball at point x , then the length of the segment decrease no more than a factor of 2




FIGURE 1. Under the conditions of Lemma 3 we have $2\|zx\| \geq \|xy\|$.

In this paper we will use the notion of the moduli of supporting convexity and smoothness, which were introduced and studied by the author in paper [19]. This technique is very convenient in questions concerning the geometry of unit ball and its supporting hyperplanes. Let us introduce the following definitions.

We say that y is *quasiorthogonal* to vector $x \in X \setminus \{o\}$ and write $y \lrcorner x$ if there exists a functional $p \in J_1(x)$ such that $\langle p, y \rangle = 0$. Note that the following conditions are equivalent:

- y is quasiorthogonal to x
- for any $\lambda \in \mathbb{R}$ the vector $x + \lambda y$ lies in the supporting hyperplane to the ball $\mathfrak{B}_{\|x\|}(o)$ at x ;
- for any $\lambda \in \mathbb{R}$ the following inequality holds $\|x + \lambda y\| \geq \|x\|$;
- x is orthogonal to y in Birkhoff–James sense ([18], Ch. 2, §1).

Let $x, y \in \partial \mathfrak{B}_1(o)$ be such that $y \lrcorner x$. By definition, put

$$\lambda_X(x, y, r) = \min \{ \lambda \in \mathbb{R} : \|x + ry - \lambda x\| = 1 \}$$

for any $r \in [0, 1]$.

Define the *modulus of supporting convexity* as

$$\lambda_X^-(r) = \inf\{\lambda_X(x, y, r) : \|x\| = \|y\| = 1, y \lhd x\}.$$

Define the *modulus of supporting smoothness* as

$$\lambda_X^+(r) = \sup\{\lambda_X(x, y, r) : \|x\| = \|y\| = 1, y \lhd x\}.$$

In paper [19] the following inequalities are proved:

$$(7) \quad \rho_X\left(\frac{r}{2}\right) \leq \lambda_X^+(r) \leq \rho_X(2r).$$

$$(8) \quad \lambda_X^-(r) \leq \delta_X(2r) \leq 1 - \sqrt{1 - r^2}.$$

$$(9) \quad 0 \leq \lambda_X^-(r) \leq \lambda_X^+(r) \leq r.$$

4. PROOF OF THEOREM 1

Auxiliary results

For the set $A \subset X$ we introduce the following function

$$\Gamma(A, \varepsilon, X) = -\inf\langle p_1 - p_2, x_1 - x_2 \rangle,$$

where the infimum is considered for all x_1, x_2 in X and p_1, p_2 in X^* that satisfy the following condition

$$(10) \quad x_i \in A, p_i \in N(A, x_i) \cap \partial\mathfrak{B}_1^*(o), \quad i = 1, 2,$$

and such that $\|x_1 - x_2\| = \varepsilon$.

The function $-\Gamma(A, \varepsilon, X)$ is the maximum value that can be on the right hand side in the hypomonotonicity inequality (2) meaning that if $A \in \Omega_N^\psi(R)$, then

$$(11) \quad -\Gamma(A, \varepsilon, X) \geq -R\psi\left(\frac{\varepsilon}{R}\right).$$

Lemma 4. *Let X be a Banach space with a Frechet differentiable norm. Let $A = X \setminus \text{int } \mathfrak{B}_1(o)$. Then*

$$(12) \quad \Gamma(A, \varepsilon, X) \asymp \lambda_X^+(\varepsilon) \asymp \rho_X(\varepsilon), \quad \varepsilon \rightarrow 0.$$

Moreover, if $\varepsilon \in [0, \frac{1}{2}]$ then the inequality

$$(13) \quad \Gamma(A, \varepsilon, X) \geq \rho_X\left(\frac{\varepsilon}{4}\right).$$

holds.

Proof.

1) The equivalence of $\lambda_X^+(\varepsilon) \asymp \rho_X(\varepsilon)$ as $\varepsilon \rightarrow 0$ is the consequence of (7).

2) Fix any $\varepsilon \in (0, \frac{1}{2}]$. As $\partial A = \partial\mathfrak{B}_1(o)$, it follows that for any $x \in \partial A$ the equality

$$(14) \quad N(A, x) \cap \partial\mathfrak{B}_1^*(o) = -J_1(x)$$

holds. Fix any x_1, x_2, p_1, p_2 , that satisfy the inclusions (10) and such that $\|x_1 - x_2\| \leq \varepsilon \leq \frac{1}{2}$. Let ℓ_1 be a line in the plane $x_1 o x_2$, supporting to the ball $\mathfrak{B}_1(o)$ at the point x_1 (its uniqueness follows from the differentiability of the norm). Let z be the projection of point x_2

on this line. Then $-\langle p_1, x_1 - x_2 \rangle = \|z - x_2\|$. From the triangle inequality it follows that $\|z - x_1\| \leq \|z - x_2\| + \|x_2 - x_1\| \leq 2\|x_2 - x_1\|$. Therefore,

$$\|z - x_2\| \leq \lambda_X^+(\|x_1 - z\|) \leq \lambda_X^+(2\|x_1 - x_2\|).$$

From the last inequality in this series we obtain that $\langle p_1, x_1 - x_2 \rangle \geq -\lambda_X^+(2\|x_1 - x_2\|)$.

Similarly, $\langle p_2, x_2 - x_1 \rangle \geq -\lambda_X^+(2\|x_1 - x_2\|)$. Summing the last two inequalities, we obtain that

$$\langle p_1 - p_2, x_1 - x_2 \rangle \geq -2\lambda_X^+(2\|x_1 - x_2\|),$$

and, so, for any $\varepsilon \in (0, \frac{1}{2}]$ we have

$$(15) \quad \Gamma(A, \varepsilon, X) \leq 2\lambda_X^+(2\varepsilon).$$

The equality (14) implies that for any x_1, x_2, p_1, p_2 , that satisfy the condition (10) the inequality

$$(16) \quad \langle p_1 - p_2, x_1 - x_2 \rangle \leq \langle p_1, x_1 - x_2 \rangle$$

holds. According to the definition of the convexity modulus, for any $\gamma > 0$ there exist $x_1, x_2 \in \partial A$; $p_1 \in N(x_1, A_1)$; $z \in H_{p_1} = \{x \in X \mid \langle p_1, x \rangle = 1\}$ such that

- (1) z is the projection of x_2 on the hyperplane H_{p_1} ;
- (2) $\|x_1 - z\| = \frac{\varepsilon}{2}$;
- (3) $\|z - x_2\| \geq \lambda_X^+(\|x_1 - z\|) - \gamma = \lambda_X^+(\frac{\varepsilon}{2}) - \gamma$.

This implies that

$$(17) \quad -\langle p_1, x_1 - x_2 \rangle = \|z - x_2\| \geq \lambda_X^+(\frac{\varepsilon}{2}) - \gamma.$$

On a unit sphere in the plane $x_1 o x_2$ we fix a point x_ε such that $\|x_1 - x_\varepsilon\| = \varepsilon$ and in the plane $x_1 o x_2$ the triangle $x_1 x_2 x_\varepsilon$ lies on one side of line $o x_1$. Lemma 3 implies that $\varepsilon = 2\|x_1 - z\| \geq \|x_1 - x_2\|$. Using the inequality (16), we obtain

$$\Gamma(A, \varepsilon, X) \geq -\langle p_1, x_1 - x_\varepsilon \rangle \geq -\langle p_1, x_1 - x_2 \rangle.$$

This and inequality (17) imply that

$$\Gamma(A, \varepsilon, X) \geq \lambda_X^+(\frac{\varepsilon}{2}) - \gamma.$$

Passing to the limit in the last inequality as $\gamma \rightarrow 0$, we obtain that

$$(18) \quad \Gamma(A, \varepsilon, X) \geq \lambda_X^+(\frac{\varepsilon}{2}).$$

The inequalities (15) and (18) imply that

$$(19) \quad 2\lambda_X^+(2\varepsilon) \geq \Gamma(A, \varepsilon, X) \geq \lambda_X^+(\frac{\varepsilon}{2})$$

Using this and (7) we obtain inequality (13).

□

Proof 2 $\Rightarrow$ 1 in Theorem 1

Fix a set $A \in \Omega_N(R)$. Theorem 4 implies that for any $x_1, x_2 \in A$ и $p_i \in N(x_i, A), i = 1, 2$ the inequality

$$(20) \quad \langle p_1 - p_2, x_1 - x_2 \rangle \geq -4R\rho_X \left(\frac{\|x_1 - x_2\|}{R} \right)$$

holds. Condition 2 implies that for some $\varepsilon > 0$ exists a positive constant k such that for any $\tau \in [0, \varepsilon]$ the inequality

$$(21) \quad \rho_X(\tau) \leq k\psi(\tau)$$

holds. The inequalities (21), (20) imply that $A \in \Omega_N^{k_1\psi}(R)$, where $k_1 = 4k$.

□

Proof 1 $\Rightarrow$ 2 in Theorem 1

Let the constant k_1 be such that $\Omega_N(1) \subset \Omega_N^{k_1\psi}(1)$. Let us show that assertion 2 holds. Consider the set $A_1 = \{x \in X \mid \|x\| \geq 1\}$. Clearly, $A_1 \in \Omega_P(1)$, and by Theorem IV the inclusion $A_1 \in \Omega_N(1)$ holds. From the definition of $\Gamma(A, \varepsilon, X)$, inequalities (13), (11) and the relations (4) we obtain for some constant $\gamma > 0$ the following series of inequalities

$$k_1\psi(\varepsilon) \geq \Gamma(A, \varepsilon, X) \geq \rho_X \left(\frac{\varepsilon}{4} \right) \geq \frac{1}{17}\rho_X(\varepsilon),$$

where $\varepsilon \in [0, \gamma]$.

□

Particularly, we have shown that the following statement holds.

Corollary 1. *In a uniformly smooth and uniformly convex Banach space X the inclusion $(X \setminus \text{int } \mathfrak{B}_1(o)) \in \Omega_N^{\frac{1}{17}\rho_X(\cdot)}(1)$ holds.*

Remark 3. *The implication 2 $\Rightarrow$ 1 in Theorem 1 is an elementary corollary of the smoothness modulus definition (see Theorem 4). But to prove the reverse implication we had to show the equivalence of smoothness and convexity moduli at zero.*

5. Proof 2 $\Rightarrow$ 1 in Theorem 2
Auxiliary results

Define the function $\omega_X : [0, +\infty) \rightarrow [0, +\infty)$ in the following way

$$\omega_X(\tau) = \frac{\rho_X(\tau)}{\tau}.$$

As $\omega_X(\cdot)$ is an increasing monotonous function, the inverse function $\omega_X^{-1}(\cdot)$ is also increasing. Note that $\omega_X(1) = \rho_X(1) \geq \rho_H(1) = \sqrt{2} - 1$, and, therefore, for any $t \in [0, 1]$ the inequality

$$(22) \quad \omega_X^{-1}((\sqrt{2} - 1)t) \leq 1,$$

holds.

The next Lemma is a generalization of Lemma 1.9.1 from G.E. Ivanov's monograph [5], in which an analogous result was obtained for Hilbert spaces.

For a set $A \subset X$ denote by $T(A)$ the set of a point x such that $P_A(x)$ is single-point. It is well known, that in a uniformly convex Banach space X for any closed set A the set $T(A)$ is dense in X (See [20]).

Lemma 5. *Let X be a uniformly smooth Banach space, A be a closed set such that the set $T(A)$ is dense in X . Let there be given $z_0 \notin A$, $z_1 \in A$ and $\varepsilon \in (0, 1)$. Then there exist $\lambda \in [0, 1]$, a point $y \in \partial A$ and $p \in N(y, A)$, $\|p\| = 1$ such that*

$$(23) \quad \|(1 - \lambda)z_0 + \lambda z_1 - y\| < \varepsilon \|z_1 - z_0\|,$$

$$(24) \quad \langle p, z_1 - z_0 \rangle < \varepsilon \|z_1 - z_0\|.$$

Proof.

Note $\delta = \frac{1}{2} \min\{\varepsilon \|z_1 - z_0\|, \rho(z_0, A)\}$. As the set A is closed and $z_0 \notin A$, we have that $\rho(z_0, A) > 0$, and, therefore $\delta > 0$. Since $2\delta \leq \rho(z_0, A)$, we obtain that $\text{int } \mathfrak{B}_{2\delta}(z_0) \subset X \setminus A$.

Define the number λ as follows

$$\lambda = \inf \left\{ t > 0 : \mathfrak{B}_\delta(z_0 + t(z_1 - z_0)) \cap A \neq \emptyset \right\}.$$

As $z_1 \in A$, we have that $\mathfrak{B}_\delta(z_1) \cap A \neq \emptyset$, and, therefore, $\lambda \leq 1$. Note that for $|t| < \frac{\delta}{\|z_0 - z_1\|}$ the inclusions

$$\mathfrak{B}_\delta(z_0 + t(z_1 - z_0)) \subset \text{int } \mathfrak{B}_{2\delta}(z_0) \subset X \setminus A,$$

hold and, therefore, $\mathfrak{B}_\delta(z_0 + t(z_1 - z_0)) \cap A = \emptyset$ as $|t| < \frac{\delta}{\|z_0 - z_1\|}$. This and the definition of number λ imply that

$$(25) \quad \lambda \geq \frac{\delta}{\|z_0 - z_1\|},$$

$$(26) \quad \mathfrak{B}_\delta(z_0 + t(z_1 - z_0)) \cap A = \emptyset, \quad \forall t \in \left(-\frac{\delta}{\|z_0 - z_1\|}, \lambda \right).$$

Define the vector $z_\lambda = (1 - \lambda)z_0 + \lambda z_1$. Since X is a uniformly smooth Banach space, there exists $\xi > 0$ such that

$$(27) \quad (\sqrt{2} + 1) \frac{\xi}{\rho_X^{-1}(\xi)} < \varepsilon$$

Define the number $\beta = \frac{\sqrt{2}-1}{2} \delta \min\{\xi, \frac{1}{2}\}$.

As the set $T(A)$ is dense in X , there exists $y_0 \in T(A)$ such that the inequality $\|z_\lambda - y_0\| < \beta$ holds. Let $\{y\} = P_A(y_0)$.

Obviously,

$$|\rho(y_0, A) - \rho(z_\lambda, A)| \leq \|y_0 - z_\lambda\| < \beta.$$

By the definition of λ we have $\rho(z_\lambda, A) = \delta$. Therefore,

$$(28) \quad \delta - \beta < \|y_0 - y\| = \rho(y_0, A) < \delta + \beta.$$

From this and the inequality $\|y_0 - z_\lambda\| < \beta$, $\beta \leq \frac{\delta}{2}$, $\delta \leq \frac{\varepsilon \|z_1 - z_0\|}{2}$ we get that

$$\|z_\lambda - y\| \leq \|y_0 - y\| + \|y_0 - z_\lambda\| < \delta + 2\beta \leq 2\delta \leq \varepsilon \|z_1 - z_0\|.$$

Thus, inequality (23) holds.

Let the functional $p \in \partial \mathfrak{B}_1^*(o)$ be dual to the vector $y_0 - y$. Lemma V implies that $p \in N(y, A)$.

If $\langle p, z_1 - z_0 \rangle \leq 0$, then inequality (24) holds. So, let us consider the case $\langle p, z_1 - z_0 \rangle > 0$. Let $\gamma = \frac{\langle p, z_1 - z_0 \rangle}{\|z_1 - z_0\|} \in (0, 1]$. We have to prove that $\gamma < \varepsilon$. Define the number

$$(29) \quad \mu = \omega_X^{-1}((\sqrt{2} - 1)\gamma) \frac{\|y - y_0\|}{\|z_1 - z_0\|}.$$

Applying to the last expression inequalities (22), (25) and (28), we obtain that

$$0 < \mu \leq \frac{\|y - y_0\|}{\|z_1 - z_0\|} \leq \frac{\delta + \beta}{\|z_1 - z_0\|} < \frac{2\delta}{\|z_1 - z_0\|} \leq \frac{\delta}{\|z_1 - z_0\|} + \lambda.$$

Thus, $-\frac{\delta}{\|z_1 - z_0\|} < \lambda - \mu < \lambda$. This and equality (26) imply that

$$\mathfrak{B}_\delta(z_0 + (\lambda - \mu)(z_1 - z_0)) \cap A = \emptyset.$$

As $y \in A$, we have that $y \notin \mathfrak{B}_\delta(z_0 + (\lambda - \mu)(z_1 - z_0)) = \mathfrak{B}_\delta(z_\lambda - \mu(z_1 - z_0))$, and so $\|z_\lambda - y - \mu(z_1 - z_0)\| > \delta$. This and the inequality $\|y_0 - z_\lambda\| < \beta$ imply that $\|y_0 - y - \mu(z_1 - z_0)\| > \delta - \beta$. Applying Lemma 1, we obtain that

$$\frac{\delta - \beta}{\|y_0 - y\|} < 1 - \mu \frac{\langle p, z_1 - z_0 \rangle}{\|y_0 - y\|} + 2\rho_X\left(\mu \frac{\|z_1 - z_0\|}{\|y_0 - y\|}\right).$$

Thus, considering inequalities (29), (28) we get that

$$\frac{\delta - \beta}{\delta + \beta} < 1 - \gamma \omega_X^{-1}((\sqrt{2} - 1)\gamma) + 2\rho_X\left(\omega_X^{-1}((\sqrt{2} - 1)\gamma)\right).$$

Note that for any $k > 0$ the equalities

$$\gamma \omega_X^{-1}(k\gamma) = \frac{\omega_X^{-1}(k\gamma) \cdot \omega_X(\omega_X^{-1}(k\gamma))}{k} = \frac{\rho_X(\omega_X^{-1}(k\gamma))}{k}$$

hold, and, therefore, we get that

$$\frac{\delta - \beta}{\delta + \beta} < 1 - \left(\frac{1}{\sqrt{2} - 1} - 2\right) \rho_X\left(\omega_X^{-1}((\sqrt{2} - 1)\gamma)\right).$$

Modifying the last inequality and using the definition of β , we obtain that

$$\rho_X\left(\omega_X^{-1}((\sqrt{2} - 1)\gamma)\right) < \frac{2(\sqrt{2} - 1)}{(3 - 2\sqrt{2})} \frac{\beta}{(\delta + \beta)} = \frac{2}{(\sqrt{2} - 1)} \frac{\beta}{(\delta + \beta)} \leq \xi,$$

and thus, considering inequality (27), we get that

$$\gamma < \frac{1}{\sqrt{2} - 1} \omega_X(\rho_X^{-1}(\xi)) = \frac{1}{\sqrt{2} - 1} \frac{\rho_X(\rho_X^{-1}(\xi))}{\rho_X^{-1}(\xi)} = (\sqrt{2} + 1) \frac{\xi}{\rho_X^{-1}(\xi)} < \varepsilon.$$

□

Proof of the main assertion

As the functions $\delta_X(\cdot)$ and $\psi(\cdot)$ are monotonous, condition 2 of theorem 2 implies that there exists a number $k > 0$ such that

$$(30) \quad R \cdot k\psi\left(\frac{t}{R}\right) \leq 2R \cdot \delta_X\left(\frac{t}{R}\right) \quad \forall t \in [0, 2R].$$

Let us prove that $\Omega_N^{k\psi}(R) \subset \Omega_N(R)$.

Suppose the contrary. Then there exists a closed set A such that $A \in \Omega_N^{k\psi}(R)$ and $A \notin \Omega_N(R)$. Then there exist a vector $x_0 \in A$, a functional $p_0 \in N(x_0, A) \cap \partial\mathfrak{B}_1^*(o)$ and the dual to it unit vector u_0 , and also the vector $z_1 \in A$ such that the inequality

$$(31) \quad \|x_0 + Ru_0 - z_1\| < R$$

holds.

According to Lemma 2, we obtain that

$$(32) \quad \langle p_0, z_1 - x_0 \rangle > 2R\delta_X \left(\frac{\|z_1 - x_0\|}{R} \right) > 0.$$

Therefore, due to the inclusion $p_0 \in N(x_0, A)$ and according to Definition 2, there exists a number $\mu \in (0, 1)$ such that the vector

$$(33) \quad z_0 = x_0 + \mu(z_1 - x_0)$$

satisfies the condition $z_0 \notin A$.

Let L be the Lipschitz constant of function $\psi(\cdot)$. Denote

$$(34) \quad \varepsilon = \frac{\mu}{(6 + 2kL)R} \left(\langle p_0, z_1 - x_0 \rangle - 2R\delta_X \left(\frac{\|z_1 - x_0\|}{R} \right) \right).$$

Inequality (32) implies that $\varepsilon > 0$.

According to Lemma 5 there exists a number $\lambda \in [0, 1]$, a vector $y \in \partial A$ and a unit functional $p \in N(y, A)$ such that

$$(35) \quad \|z_\lambda - y\| < \varepsilon \|z_1 - z_0\|,$$

$$(36) \quad \langle p, z_1 - z_0 \rangle < \varepsilon \|z_1 - z_0\|,$$

where

$$(37) \quad z_\lambda = (1 - \lambda)z_0 + \lambda z_1.$$

The definition of vector z_0 (equality (33)) implies that

$$z_1 - z_0 = (1 - \mu)(z_1 - x_0), \quad z_0 - x_0 = \mu(z_1 - x_0).$$

From this and equality (37) we get that

$$(38) \quad z_\lambda - x_0 = z_\lambda - z_0 + z_0 - x_0 = \lambda(z_1 - z_0) + z_0 - x_0 = (\lambda(1 - \mu) + \mu)(z_1 - x_0).$$

Therefore

$$\frac{z_\lambda - x_0}{\|z_\lambda - x_0\|} = \frac{z_1 - x_0}{\|z_1 - x_0\|} = \frac{z_1 - z_0}{\|z_1 - z_0\|}.$$

This and inequality (36) imply that

$$\langle p, z_\lambda - x_0 \rangle < \varepsilon \|z_\lambda - x_0\| \leq \varepsilon \|z_1 - x_0\|.$$

Thus, according to inequality (35), we have that $\langle p, y - x_0 \rangle \leq 2\varepsilon \|z_1 - x_0\|$. Inequality (31) implies that $\|z_1 - x_0\| \leq 2R$, and, therefore,

$$(39) \quad \langle p, y - x_0 \rangle \leq 4\varepsilon R.$$

As $A \in \Omega_N^{k\psi}(R)$, the inequality

$$\langle p - p_0, y - x_0 \rangle \geq -Rk\psi \left(\frac{\|y - x_0\|}{R} \right)$$

holds. From this and the inequality (39) we get that

$$(40) \quad \langle p_0, y - x_0 \rangle \leq Rk\psi\left(\frac{\|y - x_0\|}{R}\right) + 4\varepsilon R.$$

The relations (38) imply that $\|z_\lambda - x_0\| \leq \|z_1 - x_0\| < 2R$. Therefore, by inequality (35), we have that

$$\|y - x_0\| \leq \|z_\lambda - x_0\| + \varepsilon \|z_1 - z_0\| < \|z_\lambda - x_0\| + 2\varepsilon R.$$

Thus, considering the Lipschitzness of function $\psi(\cdot)$, we get that

$$Rk\psi\left(\frac{\|y - x_0\|}{R}\right) \leq Rk\psi\left(\frac{\|z_\lambda - x_0\|}{R}\right) + 2\varepsilon LkR.$$

Substituting the last inequality in (40), we obtain that

$$\langle p_0, y - x_0 \rangle \leq Rk\psi\left(\frac{\|z_\lambda - x_0\|}{R}\right) + \varepsilon R(4 + 2kL).$$

Thus, using inequality (35), we get that

$$\langle p_0, z_\lambda - x_0 \rangle < Rk\psi\left(\frac{\|z_\lambda - x_0\|}{R}\right) + \varepsilon R(6 + 2kL).$$

Denote $t = \lambda(1 - \mu) + \mu$. Then $0 < \mu \leq t \leq 1$ and, according to equality (38), we get that $z_\lambda - x_0 = t(z_1 - x_0)$. Thus,

$$t\langle p_0, z_1 - x_0 \rangle < Rk\psi\left(t\frac{\|z_1 - x_0\|}{R}\right) + \varepsilon R(6 + 2kL),$$

due to the convexity of function $\psi(\cdot)$, we obtain that

$$\langle p_0, z_1 - x_0 \rangle < Rk\frac{\psi\left(t\frac{\|z_1 - x_0\|}{R}\right)}{t} + \frac{\varepsilon R(6 + 2kL)}{t} \leq Rk\psi\left(\frac{\|z_1 - x_0\|}{R}\right) + \frac{\varepsilon R(6 + 2kL)}{\mu}.$$

Considering inequality (30), we get that

$$\langle p_0, z_1 - x_0 \rangle < 2R\delta_X\left(\frac{\|z_1 - x_0\|}{R}\right) + \frac{\varepsilon R(6 + 2kL)}{\mu},$$

which contradicts equality (34).

□

6. PROOF $1 \Rightarrow 2$ IN THEOREM 2

The proof consists in constructing examples of sets in the class $\Omega_N^{k_2\psi}(R)$ that do not belong to the class $\Omega_N(R)$, and consists in number of steps. The main idea is building in the plane $L \subset X$ a set A_1 that belongs to the class $\Omega_N^{k\psi}(R)$ in space L . Then, with the help of some technical lemmas, the set A_1 extends to the set A , which already is in the class $\Omega_N^{K\psi}(K_1R)$ in X , where the constants k, K, K_1 do not depend on the initially chosen plane L and space X . Using the fact that $\delta_X(t) \asymp \lambda_X^-(t) = o(\psi(t))$ at zero, we chose such vectors that their existence contradicts the N -supporting condition.

Auxiliary results

Let us first describe the functions ψ from the class $\mathfrak{M}$, such that there exists a normed space

whose smoothness modulus is equivalent to the function ψ . In paper [11] the following theorem is proved.

Theorem VI. *For the function $N : [0, +\infty) \rightarrow [0, +\infty)$, that satisfies the Figiel condition and such that $N(0) = 0$ there exists a two-dimensional space X_2 whose smoothness modulus $\rho_{X_2}(t)$ is equivalent to $N(t)$ at zero.*

The Day-Nordlander Theorem (see [18]) implies that if for function ψ there exists a Banach space with the smoothness modulus, equivalent to ψ at zero, then $t^2 = O(\psi(t))$. However, the function $\psi(\cdot)$ from $\mathfrak{M}$ such that $t^2 = O(\psi(t))$ may not satisfy the Figiel condition (see (3)), but the following Lemma holds.

Lemma 6. *For any function $\psi \in \mathfrak{M}$ such that $t^2 = o(\psi(t))$ at zero there exists a function $\psi_1(\cdot) : [0, +\infty) \rightarrow [0, +\infty)$ that satisfies the Figiel condition and such that $t^2 = o(\psi_1(t))$ at zero and $\psi_1(t) = o(\psi(t))$ at zero.*

Proof.

Clearly, it is sufficient to examine the case $\psi(t) = o(t)$ at zero, otherwise $\psi_1(t) = t^{\frac{3}{2}}$.

Consider the function $h(t) = \frac{t^2}{\psi(t)}$ on the interval $[0, 1]$. The function is continuous on this interval and for some constant $k > 0$ the inequality $h(t) \geq kt$ holds.

As the set $K = \{(t, y) \mid t \in [0, 1], 0 \leq y \leq h(t)\}$ is compact, its convex hull $\text{co } K$ is also a compact set.

Define a continuous function $\gamma(t) = \max\{y \in \mathbb{R} \mid (t, y) \in \text{co } K\}$, $t \in [0, 1]$. The function $\gamma(\cdot)$ is concave on the interval $[0, 1]$, $\gamma(0) = 0$. Therefore, in some neighborhood of zero $\gamma(\cdot)$ strictly increases.

Define the function $\psi_1(t) = \frac{t^2}{\sqrt{\gamma(t)}}$. As $\frac{\psi_1(t)}{\psi(t)} = \frac{h(t)}{\sqrt{\gamma(t)}} \leq \sqrt{\gamma(t)} \rightarrow 0$ and $\frac{t^2}{\psi_1(t)} = \sqrt{\gamma(t)}$ approaches zero monotonically as $t \rightarrow 0$, then the function $\psi_1(\cdot)$ satisfies the required conditions.

□

Now let us prove the technical lemmas, with the help of which we will extend the set A_1 , which lies in the class $\Omega_N^{k\psi}(R)$ on a plane, to the set A , which already lies in the class $\Omega_N^{K\psi}(K_1R)$ in the space, which contains the plane, where the constants k, K, K_1 do not depend on the choice of the initial plane and the space containing it.

If the Banach space can be represented as the direct sum of two of its closed subspaces $Z = Z_1 \oplus Z_2$, then any vector $z \in Z$ is uniquely expressible as the sum of two vectors $z_1 \in Z_1$ and $z_2 \in Z_2$, and in this case we will write $z = (z_1, z_2)$. Hereafter, when we speak about the representation of a Banach space as the direct sum of its subspaces we will consider that the subspaces are closed.

Lemma 7. *In the Banach space X fix any $x \in \partial\mathfrak{B}_1(o)$, $p \in J_1(x)$, $y \in \text{Ker } p \cap \partial\mathfrak{B}_1(o)$ (i.e. $y \perp x$). Denote $L = \text{Lin } \{x, y\}$, $l = \text{Lin } \{y\}$. Then there exists a projector $P : X \rightarrow L$ such that $P(\text{Ker } p) = l$ and $\|P\| \leq 3$.*

Proof.

If $\dim X = 2$, then the identity transformation satisfies the required condition.

Let $\dim X > 2$. According to the paper [13], in the subspace $\text{Ker } p$ of co-dimension 1 there exists a projector $P_1 : \text{Ker } p \rightarrow l$ with unit norm. Clearly, X is the direct sum of the subspaces $\text{Lin } x$ and $\text{Ker } p$.

Define the projector $P : X \rightarrow L$ in the following way

$$Pz = P(z_1, z_2) = (z_1, P_1 z_2), \quad \forall z = (z_1, z_2) \in X, \quad z_1 \in \text{Lin } x, \quad z_2 \in \text{Ker } p.$$

Let us estimate its norm. Fix an arbitrary vector $u = (u_1, u_2) \in \partial \mathfrak{B}_1(o)$, where $u_1 \in \text{Lin } x$, $u_2 \in \text{Ker } p$. As $u_2 \perp u_1$, then $\|u_1\| \leq \|u_1 + u_2\| = \|u\| = 1$. From this and the triangle inequality we get that $\|u_2\| \leq \|u_1 + u_2\| + \|u_1\| \leq 2$. Therefore,

$$\|Pu\| = \|(u_1, P_1 u_2)\| = \|u_1 + P_1 u_2\| \leq \|u_1\| + \|P_1 u_2\| \leq \|u_1\| + \|u_2\| \leq 3.$$

□

Lemma 8. *Let X be a linear vector space with two equivalent norms $\|\cdot\|_1$ and $\|\cdot\|_2$:*

$$(41) \quad \|x\|_1 \leq \|x\|_2 \leq c_1 \|x\|_1.$$

Denote $X_1 = (X, \|\cdot\|_1)$, $X_2 = (X, \|\cdot\|_2)$. Let $A = X_2 \setminus \text{int } \mathfrak{B}_1(o)$ and $A \in \Omega_N^\psi(R)$ in the space X_2 , then $A \in \Omega_N^{c_1^2\psi}\left(\frac{R}{c_1}\right)$ in the space X_1 .

Proof.

According to Remark 1, at every point of set A the normal cone does not change, but the norms of the corresponding functionals may change. Fix arbitrary x_1, x_2, p_1^1, p_2^1 , which satisfy the condition (10) for the space X_1 . For $j = 1, 2$ denote $p_j^2 = \frac{p_j^1}{\|p_j^1\|_2}$. As $\|p_j^1\|_1 = 1$, then by inequalities (41) we have $\frac{1}{c_1} \leq \|p_j^1\|_2 \leq 1$. As the set A is a complement of a convex set, then $\langle p_1^1, x_1 - x_2 \rangle \leq 0$. Therefore,

$$\langle p_1^2, x_1 - x_2 \rangle \geq \langle p_1^1, x_1 - x_2 \rangle \geq c_1 \langle p_1^2, x_1 - x_2 \rangle.$$

Similarly, $\langle p_2^2, x_2 - x_1 \rangle \geq \langle p_2^1, x_2 - x_1 \rangle \geq c_1 \langle p_2^2, x_2 - x_1 \rangle$. Summing the last two series of inequalities, we obtain that

$$(42) \quad \langle p_1^2 - p_2^2, x_1 - x_2 \rangle \geq \langle p_1^1 - p_2^1, x_1 - x_2 \rangle \geq c_1 \langle p_1^2 - p_2^2, x_1 - x_2 \rangle.$$

From the fact that $A \in \Omega_N^\psi(R)$ in the space X_2 and from the inequalities (41) we have that

$$c_1 \langle p_1^2 - p_2^2, x_1 - x_2 \rangle \geq -c_1 R \psi\left(\frac{\|x_1 - x_2\|_2}{R}\right) \geq -c_1 R \psi\left(\frac{c_1 \|x_1 - x_2\|_1}{R}\right).$$

This and the last inequality in sequence (42) imply that

$$\langle p_1^1 - p_2^1, x_1 - x_2 \rangle \geq -c_1^2 R \psi\left(\frac{c_1 \|x_1 - x_2\|_1}{R}\right).$$

And, thus, $A \in \Omega_N^{c_1^2\psi}\left(\frac{R}{c_1}\right)$ in the space X_1 .

□

Let $X = X_1 \oplus X_2$, and, besides, there exists a projector $P : X \rightarrow X_1$ with the norm $\|P\| < +\infty$ and $X_2 = \text{Ker } P$. Then $X^* = X_1^\perp \oplus X_2^\perp$, where the spaces $X_1^\perp, X_2^\perp$ are the right annihilators to the spaces $X_2 \subset X$ and $X_1 \subset X$ accordingly. Continuing any functional $p \in X_1^*$ with zero on the space X_2 , we obtain a natural isomorphism $X_1^* = X_2^\perp$. Similarly, $X_2^* = X_1^\perp$.

Lemma 9. *Let $X = X_1 \oplus X_2$, and, besides, there exists a projector $P : X \rightarrow X_1$ with the norm $\|P\| < +\infty$ and $X_2 = \text{Ker } P$. Let the set $A_1 \subset X_1$. Define the set $A = \{x \in X \mid Px \in A_1\}$. Then $N(x, A) = (N(Px, A_1), o) = \{(p, o) \mid p \in N(Px, A_1)\}$ for any $x \in A$.*

Proof.

Fix an arbitrary vector $x = (x_1, x_2) \in A$.

1) Let $p = (p_1, p_2) \in N(x, A)$. Let us prove that $p_2 = o$. Suppose that $p_2 \neq o$. Then there exists a non-zero vector $v = (o, v_2)$ such that $\langle p, v \rangle = \langle p_2, v_2 \rangle = k \neq 0$. According to the definition of set A for any $\lambda \in \mathbb{R}$ the vector $x + \lambda v$ lies in A . But then $\langle p, \lambda v \rangle = \lambda k$ is a linear in λ function and, therefore, $p \notin N(x, A)$. Contradiction.

2) Let $p = (p_1, o) \in N(x, A)$. Let us show that $p_1 \in N(Px, A_1) = N(x_1, A_1)$. Suppose the contrary, that $p_1 \notin N(x_1, A_1)$. Then there exists $\varepsilon > 0$ such that for any $\delta > 0$ there exists a vector $y_1^\delta \in A_1$ such that $\|y_1^\delta - x_1\| \leq \delta$ and also $\langle p_1, y_1^\delta - x_1 \rangle \geq \varepsilon \|y_1^\delta - x_1\|$. Define $y_\delta = (y_1^\delta, x_2)$. Clearly, $y_\delta \in A$. Then from the last inequality we have that

$$\langle p, y_\delta - x \rangle = \langle (p_1, o), (y_1^\delta - x_1, o) \rangle = \langle p_1, y_1^\delta - x_1 \rangle \geq \varepsilon \|y_1^\delta - x_1\| = \varepsilon \|y_\delta - x\|.$$

Therefore, $p \notin N(x, A)$. Contradiction.

3) Let $p_1 \in N(x_1, A_1)$. Let us show that the inclusion $p = (p_1, o) \in N(x, A)$ holds. Fix $\varepsilon > 0$. According to the definition of the normal cone, for $\frac{\varepsilon}{\|P\|}$ there exists $\delta > 0$ such that for any vector $d_1 \in \mathfrak{B}_\delta(x_1) \cap A_1$ the inequality $\langle p_1, d_1 - x_1 \rangle \leq \frac{\varepsilon}{\|P\|} \|d_1 - x_1\|$ holds. For any $y = (y_1, y_2) \in \mathfrak{B}_{\frac{\delta}{\|P\|}}(x) \cap A$ we have $\|P(y - x)\| = \|y_1 - x_1\|_1 \leq \|P\| \|y - x\| \leq \delta$. This and the previous inequality imply that

$$\langle p, y - x \rangle = \langle p_1, y_1 - x_1 \rangle \leq \frac{\varepsilon}{\|P\|} \|y_1 - x_1\| \leq \varepsilon \|y - x\|.$$

□

Lemma 10. *If, additionally, in the assertion of Lemma 9 the set A_1 is a complement to a convex set in X_1 and $A_1 \in \Omega_N^\psi(R)$ in X_1 , then $A \in \Omega_N^{\|P\|\psi}\left(\frac{R}{\|P\|}\right)$ in the space X .*

Proof.

Through $\|\cdot\|_1$ denote the norm of space X_1^* . Clearly, for $p = (p_1, o) \in X^*$ the following inequality

$$(43) \quad \|p_1\|_1 \leq \|p\|$$

holds. Let $x = (x_1, x_2), y = (y_1, y_2) \in A$, $p^x \in N(x, A) \cap \partial \mathfrak{B}_1^*(o)$, $p^y \in N(y, A) \cap \partial \mathfrak{B}_1^*(o)$ and the following inequality holds $\|x_1 - y_1\| \leq \varepsilon$, where ε is taken from definitions of the class $\Omega_N^\psi(R)$. Lemma 9 implies that $p^x = (p_1^x, o)$, $p^y = (p_1^y, o)$ and $p_1^x \in N(x_1, A_1)$, $p_1^y \in N(y_1, A_1)$ in the space X_1 . As A_1 is a complement to a convex set in the space X_1 , then the set A is a complement to a convex set in X . Then $\langle p^x, x - y \rangle \leq 0$. From this and the inequalities (43) we get that

$$\langle p^x, x - y \rangle = \langle p_1^x, x_1 - y_1 \rangle \geq \left\langle \frac{p_1^x}{\|p_1^x\|_1}, x_1 - y_1 \right\rangle.$$

Similarly, $\langle p^y, y - x \rangle \geq \left\langle \frac{p_1^y}{\|p_1^y\|_1}, y_1 - x_1 \right\rangle$. Summing the last two inequalities, we obtain that

$$\begin{aligned} \langle p^x - p^y, x - y \rangle &\geq \left\langle \frac{p_1^x}{\|p_1^x\|_1} - \frac{p_1^y}{\|p_1^y\|_1}, x_1 - y_1 \right\rangle \geq -R\psi\left(\frac{\|x_1 - y_1\|}{R}\right) \geq \\ &\geq -R\psi\left(\frac{\|P\| \|x - y\|}{R}\right). \end{aligned}$$

□

In the next lemma there are constructed examples of sets that belong to the class $\Omega_N^{k_2\psi}(R)$ and do not belong to the class $\Omega_N(R)$. To prove it we will need some additional results from the geometry of convex sets and Banach spaces.

Definition 8. For a convex set $K \subset \mathbb{R}^n$ the John ellipsoid is called the maximum volume ellipsoid that is contained within K .

It is known that for any convex $K \subset \mathbb{R}^n$ such an ellipsoid exists and is unique. Moreover, the following theorem holds (see [14]).

Theorem VII. Let B_E be the John ellipsoid of the unit ball B_n in the space X_n , $\dim X_n = n$. Then the following inclusions

$$(44) \quad B_E \subset B_n \subset \sqrt{n}B_E$$

hold.

According to paper [15], the convexity modulus of an arbitrary Banach space X satisfies the relation

$$(45) \quad \frac{\delta_X(s)}{s^2} \leq L \frac{\delta_X(t)}{t^2} \quad \forall 0 < s \leq t \leq 2$$

for some constant $0 < L < 4$.

Lemma 11. Let there be given a function $\psi(\cdot) \in \mathfrak{M}$ such that $t^2 = O(\psi(t))$ as $t \rightarrow 0$. Let the convexity modulus $\delta_X(\cdot)$ in the Banach space X satisfy the equality $\delta_X(t) = o(\psi(t))$ as $t \rightarrow 0$. Then there does not exist a constant $k_2 > 0$ such that the inclusion $\Omega_N^{k_2\psi}(R) \subset \Omega_N(R)$ holds for any $R > 0$.

Proof.

Suppose the contrary, that there exists a constant $k_2 > 0$ such that $\Omega_N^{k_2\psi}(R) \subset \Omega_N(R)$. As the condition of Lemma 11 implies that $\delta_X(\varepsilon) = o(\psi(\varepsilon))$ as $\varepsilon \rightarrow 0$, it is sufficient to consider two cases of relations between the functions $t^2, \delta_X(t), \psi(t)$ at zero:

- 1) $t^2 = o(\psi(t))$ as $t \rightarrow 0$;
- 2) $\delta_X(t) = o(t^2)$ as $t \rightarrow 0$.

Indeed, if $t^2 = O(\psi(t))$ and $t^2 \neq o(\psi(t))$ as $t \rightarrow 0$, then $\liminf_{t \rightarrow 0} \frac{\psi(t)}{t^2} = c \in (0, +\infty)$. This and the equality $\delta_X(t) = o(\psi(t))$ as $t \rightarrow 0$ imply that $\liminf_{t \rightarrow 0} \frac{\delta_X(t)}{t^2} = 0$. From this and inequality (45) we obtain that $\delta_X(t) = o(t^2)$ as $t \rightarrow 0$.

We now pass to the consideration of the two cases, mentioned above.

The case $t^2 = o(\psi(t))$ as $t \rightarrow 0$.

Lemma 6 and Theorem VI imply that in a two-dimensional linear space Y there exists a norm $\|\cdot\|_s$ such that the smoothness modulus $\rho_{X_s}(\cdot)$ of the space $X_s = (Y, \|\cdot\|_s)$ satisfies the conditions $t^2 = o(\rho_{X_s}(t))$ and $\rho_{X_s}(t) = o(\psi(t))$ as $t \rightarrow 0$.

Therefore, there exists a constant γ_0 such that for any $\gamma \in (0, \gamma_0]$ the inequality

$$(46) \quad \frac{\rho_{X_s}\left(\frac{\gamma}{4}\right)}{16\gamma} \geq 51\gamma$$

holds.

Fix $0 < \varepsilon < \min \left\{ \frac{1}{2}, \gamma_0 \right\}$. By the definition of the modulus of supporting convexity and (8), in space X there exist vectors $u, v \in \partial \mathfrak{B}_1(o)$ such that $v \lhd u$ and

$$(47) \quad \lambda_X(u, v, \varepsilon) \leq 2(1 - \sqrt{1 - \varepsilon^2}) < 2\varepsilon^2.$$

Let $p \in -J_1(u)$ be such that $\langle p, v \rangle = 0$.

Consider the plane $X_2 = \text{Lin} \{u, v\}$. Note that the restriction of the functional p on X_2 does not change its norm. Denote this restriction by p_1 . Through B_2 denote the unit ball in this space with the norm induced by the space X . Consider the John ellipsoid (ellipse) B_E for $\frac{B_2}{\sqrt{2}}$. Denote the norm, generated by the set B_E as a unit ball through $\|\cdot\|_E$. Inclusion (44) implies that for any $x \in X_2$ the inequality $2\|x\| \geq \|x\|_E \geq \sqrt{2}\|x\|$ holds. There exists an affine transformation that transforms the John ellipsoid (ellipse) of the space X_s into the ellipsoid B_E . Replacing the space X_s by the isometrically isomorphic to it space, which is obtained from the affine transformation, we consider that the set B_E is the John ellipsoid of the unit ball B_s in the space X_s . Moreover, the inequality (46) holds. Therefore, the norms $\|\cdot\|_E, \|\cdot\|, \|\cdot\|_s$ in a two-dimensional linear space are related by

$$(48) \quad \|x\| \leq \frac{\|x\|_E}{\sqrt{2}} \leq \|x\|_s \leq \|x\|_E \leq 2\|x\|.$$

Denote $\gamma = \frac{\varepsilon}{25}$. In the space X_s , according to the definition of $\Gamma(X_s \setminus \text{int } B_s, \gamma, X_s)$ and the inequality (13), there exist d and f from the unit ball in X_s such that $\|d - f\|_s = \gamma$ and for $q \in -J_1(f)$ the inequality

$$\langle q, d - f \rangle \geq \frac{1}{2} \cdot \frac{1}{2} \Gamma(X_s \setminus \text{int } B_s, \gamma, X_s) \geq \frac{1}{4} \rho_{X_s} \left(\frac{\gamma}{4} \right)$$

holds.

Executing the affine transformation K , which, if necessary, consists in the symmetry with respect to the main axis of the ellipse B_E and rotation by some angle ϕ_0 with respect to the ellipse B_E (i.e. K is such an affine transformation of the plane X_2 that the ellipse B_E transforms into itself), we can achieve that

- (1) The line l , supporting to $K(B_s)$ at point Kf , is parallel to $\text{Ker } p = \{\lambda v \mid \lambda \in \mathbb{R}\}$.
- (2) The projection Kd on l along u lies in ray $\{Kf + \lambda v \mid \lambda \geq 0\}$.

Denote $A_1 = X_s \setminus \text{int } K(B_s)$, $f_1 = Kf$, $d_1 = Kd$. Using the fact that the linear functional on a two-dimensional space is defined up to a multiplicative constant by the line that is the kernel of the functional, the functional $q_1 \in -J_1(f_1)$ (unique due to the smoothness of B_s) is codirectional with the functional p_1 (see Fig. 2).

Let $q_1 = kp_1$. Then, as the transformation K is affine, we obtain that

$$(49) \quad \langle kp_1, d_1 - f_1 \rangle \geq \frac{1}{4} \rho_{X_s} \left(\frac{\gamma}{4} \right).$$

On the other hand, it is clear that the norm $\|\cdot\|'_s$ of the space X'_s , generated by the set $K(B_s)$ (i.e. we consider $K(B_s)$ to be the unit ball), satisfies the following series of inequalities

$$(50) \quad \|x\| \leq \frac{\|x\|_E}{\sqrt{2}} \leq \|x\|'_s \leq \|x\|_E \leq 2\|x\|.$$




FIGURE 2. To the proof of Lemma 11. In the Lemma notations:

- with a black dotted line the boundaries of the ellipses $= \frac{B_E}{\sqrt{2}}, B_E, \sqrt{2}B_E$ are shown;
- with a blue line the boundaries of the balls $B_2, \frac{B_2}{2}$ are depicted;
- with brown the boundaries of balls B_s, KB_s are depicted;
- ℓ_1 is the ray $\{u + \lambda v | \lambda \geq 0\}$ (part of the supporting line $\text{Ker } p_1 + u$);
- l' is the preimage of line l (the supporting line to the ball B_s at point f).

This, particularly, implies that $1 \leq k \leq 2$. As the right side of the inequality (49) is positive, we obtain that

$$(51) \quad \langle p_1, d_1 - f_1 \rangle \geq \frac{1}{8} \rho_{X'_s} \left(\frac{\gamma}{4} \right).$$

According to Lemma 7, in X there exists a projector $P : X \rightarrow X_2$ with the norm not greater than 3, that projects $\text{Ker } p$ on $\text{Ker } p_1$. Define the set $A = \{x \in X | Px \in A_1\}$. According to Corollary 1, $A_1 \in \Omega_N^{\frac{1}{17}\rho_{X'_s}(\cdot)}(1)$ in the space X'_s . Then Lemma 8 and the series of inequalities (50) imply that $A_1 \in \Omega_N^{\frac{4}{17}\rho_{X'_s}(\cdot)}(\frac{1}{2})$ in the space X_2 . This, according to Lemma 10, implies that the inclusion $A \in \Omega_N^{\frac{12}{17}\rho_{X'_s}(\cdot)}(\frac{1}{6})$ in space X holds. But by construction we have that $\rho_{X'_s}(t) = \rho_{X_s}(t) = o(\psi(t))$ as $t \rightarrow 0$. Therefore, for sufficiently small t the inequality $\frac{12}{17}\rho_{X'_s}(6t) \leq k_2\psi(6t)$ holds, which implies that $A \in \Omega_N^{k_2\psi(\cdot)}(\frac{1}{6})$. According to the assumption made, we obtain that $A \in \Omega_N(\frac{1}{6})$.

By construction for the point $f_1 \in X_2 \subset X$ the equality $p = (p_1, o)$ holds, and, therefore, $p = N(f_1, A) \cap \partial \mathfrak{B}_1^*(o)$. Let $D = \mathfrak{B}_{\frac{1}{6}}(f_1 - \frac{1}{6}u)$. According to the definition of the class $\Omega_N(\frac{1}{6})$, the following relation is satisfied

$$(52) \quad \text{int } D \cap A = \emptyset.$$

The inequality (47) implies that $\frac{\lambda_X(u, v, \varepsilon)}{6} < \frac{\varepsilon^2}{3} < \frac{1}{6}$. By the definition of $\lambda_X(u, v, \varepsilon)$ and the choice of vectors u and v we have that $z_1 = f_1 + \frac{\varepsilon}{6}v - \frac{\varepsilon^2}{3}u \in \text{int } D$. In the space X_2 consider the projection h_1 of the point d_1 on the line l . By the uniform convexity and uniform smoothness

of the space X_2 the segment h_1d_1 is parallel to u . On the other hand, as the vector $d_1 \in X$ lies in A , then from (52) we obtain that $d_1 \notin \text{int } D$. Therefore, the segment f_1d_1 intersects the sphere ∂D in two points. Then, by construction, the following series of inequalities holds $\|h_1 - f_1\| \leq \|f_1 - d_1\| + \|d_1 - h_1\| \leq 2\|f_1 - d_1\| \leq 2\|f_1 - d_1\|'_s = 2\gamma < \frac{\varepsilon}{6} < \frac{1}{6}$. From this, (9) and considering that the segment $h_1d_1 \cap \text{int } D = \emptyset$, $h_1d_1 \parallel u$, the point h_1 lies on the supporting line to the ball D of radius $\frac{1}{6}$ at the point f_1 , we get that

$$\|d_1 - h_1\| \leq \frac{1}{6} \lambda_{X_2} \left(f_1, \frac{h_1 - f_1}{\|h_1 - f_1\|}, 6\|h_1 - f_1\| \right) \leq \|h_1 - f_1\|,$$

and, thus, $\|f_1 - d_1\| \leq 2\|h_1 - f_1\|$. From this, the choice of f_1, d_1 and the inequality (50), we obtain that

$$(53) \quad 0 < \frac{\gamma}{2} = \frac{\|f_1 - d_1\|'_s}{2} \leq \|f_1 - d_1\| \leq 2\|h_1 - f_1\| \leq 4\gamma < \frac{\varepsilon}{6}.$$

But $d_1 = f_1 + \|h_1 - f_1\|v - \langle p_1, d_1 - f_1 \rangle u = f_1 + \|h_1 - f_1\|v - \langle p, d_1 - f_1 \rangle u$. The inequalities (53), (51), (46) and the choice of constant γ imply that

$$(54) \quad \frac{\langle p, d_1 - f_1 \rangle}{\|h_1 - f_1\|} \geq \frac{\rho_{X_s}(\frac{2}{4})}{16\gamma} \geq 51\gamma > 2\varepsilon.$$

From $\|f_1 - h_1\| < \frac{\varepsilon}{6}$ and inequality (54) we obtain that the segments $d_1h_1 \subset A$ and $f_1z_1 \subset D$ intersect. Denote by y_1 their intersection point. Clearly, $y_1 \in \text{int } D \cap A$ (see Fig. 3). This contradicts relation (52).




FIGURE 3. To the proof of Lemma 11. In the notations of the lemma:

- $o' = f_1 - \frac{1}{6}u$, $b = o' + \frac{1}{6}v$, $a_1 = f_1 + \frac{1}{6}v$, $g_1 = f_1 + \frac{\varepsilon}{6}v$;
- as $\frac{\lambda_X(u, v, \varepsilon)}{6} < \|z_1 - g_1\| < \frac{1}{6}$, then z_1 lies inside the parallelogram $o'ba_1f_1$ and inside D ;
- similarity implies that $\frac{\|z_1 - g_1\|}{\|f_1 - g_1\|} = \frac{\|y_1 - h_1\|}{\|f_1 - h_1\|} < \frac{\|d_1 - h_1\|}{\|f_1 - h_1\|}$.

The case $\delta_X(t) = o(t^2)$ as $t \rightarrow 0$.

The consideration of this case does not differ a lot from the consideration of the previous one. The difference consists in the specific choice of constants. All the notations coincide with the

notations in the previous case, if not mentioned otherwise. Let $k_3 > 0$ be such that $\psi(t) \geq k_3 t^2$ in some neighborhood of zero. Note that if $t \in [0, 1]$ the inequalities

$$(55) \quad \frac{t^2}{4} \leq \rho_H(t) = \sqrt{1+t^2} - 1 \leq t^2$$

hold.

Let $R_1 = \frac{17}{72}k_2k_3$. By assumption we have that $\delta_X(t) = o(t^2)$ as $t \rightarrow 0$. Fix $\gamma_0 \in (0, 1)$ such that for any $t \in (0, \gamma_0)$ the inequality $\delta_X(t) < \frac{1}{3} \cdot \frac{R_1}{2^{12}}t^2$ holds.

By the definition of the modulus of supporting convexity and (8) for

$$\varepsilon \in \left(0, \min\left\{\frac{\gamma_0}{2}, \frac{1}{\sqrt{R_1}}\right\}\right)$$

there exist $u, v \in \partial \mathfrak{B}_1(o)$ in the space X such that $v^\top u$ and

$$(56) \quad \lambda_X(u, v, \varepsilon) \leq 2\delta_X(2\varepsilon) < \frac{1}{3} \cdot \frac{R_1}{512}\varepsilon^2 < 1.$$

The space X_2 and the functionals p, p_1 are defined similarly to the previous case. We consider the unit ball in X'_s to be the set B_E . Fix $\gamma = \frac{R_1\varepsilon}{3}$. Similarly to the previous case, we take the points f_1, d_1, h_1 , and construct the set A . As in the previous case, $A \in \Omega_N^{\frac{12}{17}\rho_{X'_s}(\cdot)}(\frac{1}{6})$. As the ball in X'_s is Euclidean and considering the left inequality in series (55) we have that

$$(57) \quad \langle p_1, d_1 - f_1 \rangle \geq \frac{1}{4}\rho_{X'_s}\left(\frac{\gamma}{4}\right) \geq \frac{1}{256}\gamma^2.$$

From the fact that the norm in X'_s is Euclidean, the right hand side of inequality (55) and the choice of constants k_3 и R_1 we obtain that

$$R_1k_2\psi\left(\frac{t}{R_1}\right) \geq \frac{k_2k_3t^2}{R_1} = \frac{72t^2}{17} \geq \frac{1}{6} \cdot \frac{12}{17}\rho_{X'_s}(6t).$$

Thus, $A \in \Omega_N^{k_2\psi(\cdot)}(R_1)$. The assumption implies that $A \in \Omega_N(R_1)$.

Let $D = \mathfrak{B}_{R_1}(f_1 - R_1u)$. Inequality (56) and the choice of constant ε imply that $R_1\lambda_X(u, v, \varepsilon) < \frac{1}{3} \cdot \frac{R_1^2}{512}\varepsilon^2 < R_1$. By the definition of $\lambda_X(u, v, \varepsilon)$ and the choice of vectors u and v we get that $z_1 = f_1 + R_1\varepsilon v - \frac{1}{3} \cdot \frac{R_1^2}{512}\varepsilon^2 u \in \text{int } D$.

Similarly to the previous case, the estimations of the norm $h_1 - f_1$ are made

$$(58) \quad \frac{\gamma}{4} \leq \|h_1 - f_1\| \leq 2\gamma < R_1\varepsilon.$$

Furthermore, $d_1 = f_1 + \|h_1 - f_1\|v - \langle p, d_1 - f_1 \rangle u$. Inequalities (58), (57) and the choice of constant γ imply that

$$(59) \quad \frac{\langle p, d_1 - f_1 \rangle}{\|h_1 - f_1\|} \geq \frac{1}{512}\gamma = \frac{1}{3} \cdot \frac{1}{512}R_1\varepsilon.$$

By the inequality $\|f_1 - h_1\| < R_1\varepsilon$ and inequality (59) we get that the segments $d_1h_1 \subset A$ and $f_1z_1 \subset D$ intersect. Denote their intersection point by y_1 . Clearly, $y_1 \in \text{int } D \cap A$. This contradicts the relation (52).

□

Proof of the main theorem.

Let us show that the proof $1 \Rightarrow 2$ in Theorem 2 follows from Lemma 11.

Inequalities (45), (3) imply that if for the function $\psi \in \mathfrak{M}_2$ the relation $\psi(\varepsilon) \neq O(\delta_X(\varepsilon))$ holds as $\varepsilon \rightarrow 0$, then $\delta_X(\varepsilon) = o(\psi(\varepsilon))$ as $\varepsilon \rightarrow 0$.

□

6.1. Proof of Theorem 3.

Lemma 12. *Let X be a uniformly smooth and uniformly convex Banach space. Let there be given a function $\psi(\cdot) \in \mathfrak{M}$ such that $t^2 = O(\psi(t))$ as $t \rightarrow 0$. If there exists a constant $k_2 > 0$ such that the inclusion $\Omega_N^{k_2\psi}(R) \subset \Omega_N(R)$ holds, then $\delta_X(\varepsilon) \asymp \varepsilon^2$ at zero and $\liminf_{t \rightarrow 0} \frac{\psi(t)}{t^2} = c \in (0, +\infty)$.*

Proof.

Applying Lemma 11, we get that $\delta_X(\varepsilon) \neq o(\psi(\varepsilon))$ as $\varepsilon \rightarrow 0$. But for the function $\psi(\cdot) \in \mathfrak{M}$ such that $t^2 = O(\psi(t))$ as $t \rightarrow 0$, the relation $\delta_X(\varepsilon) \neq o(\psi(\varepsilon))$ as $\varepsilon \rightarrow 0$ may hold if and only if $\delta_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$ and $\liminf_{t \rightarrow 0} \frac{\psi(t)}{t^2} = c \in (0, +\infty)$.

□

Proof of Theorem 3.

Applying Theorem 1, we obtain that in some neighborhood of zero for some constant $k > 0$ the inequality

$$(60) \quad \psi(\varepsilon) \geq k\rho_X(\varepsilon)$$

holds. This and the Day-Nordlander theorem (see [18]) imply that $\varepsilon^2 = O(\psi(\varepsilon))$ as $\varepsilon \rightarrow 0$. Now, applying Lemma 12, we obtain that $\delta_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$ and $\liminf_{t \rightarrow 0} \frac{\psi(t)}{t^2} = c \in (0, +\infty)$.

This, inequality (60) and the fact that the smoothness modulus satisfies the Figiel condition (see Remark 2) imply that $\rho_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$.

Since $\delta_X(\varepsilon) \asymp \rho_X(\varepsilon) \asymp \varepsilon^2$ as $\varepsilon \rightarrow 0$, X is isomorphic to a Hilbert space (see [16], §5.5.6).

□

REFERENCES

- [1] F. H. Clarke, R. J. Stern, and P. R. Wolenski. Proximal smoothness and lower- C^2 property. *J. Convex Analysis*, 2(1):117–144, 1995.
- [2] R.A. Poliquin, R.T. Rockafellar. Prox-regular functions in variational analysis. *Trans. Amer. Math. Soc.*, 368:1805–1838, 1996.
- [3] R.A. Poliquin, R.T. Rockafellar, and L. Thibault. Local differentiability of distance functions. *Trans. Amer. Math. Soc.*, 352(11):5231–5249, 2000.
- [4] J.-P. Vial. Strong and weak convexity of sets and functions. *Math. Ops. Res.*, 8(2):231–259, 1983.
- [5] G. E. Ivanov. *Weakly convex sets and functions. Theory and applications*. Fizmatlit (in Russian), 2006.
- [6] F. Bernard, L. Thibault, and N. Zlateva. Characterizations of prox-regular sets in uniformly convex banach spaces. *J. Convex Anal.*, 13:525–559, 2006.
- [7] F. Bernard, L. Thibault, and N. Zlateva. Prox-regular sets and epigraphs in uniformly convex Banach spaces: Various regularities and other properties. *Trans. Amer. Math. Soc.*, 363:2211–2247, 2011.
- [8] G. E. Ivanov, G. M. Ivanov. Relationship between supporting conditions in Banach spaces. *Proceedings of MIPT*, 3(1):74–80, 2011.
- [9] M. V. Balashov and G. E. Ivanov. Weakly convex and proximally smooth sets in banach spaces. *Izv. RAN. Ser. Mat.*, 73(3):23–66, 2009.
- [10] J. Lindenstrauss and L. Tzafriri. *Classical Banach Spaces II*. Springer-Verlag, 1979.

- [11] A. J. Guirao, M. Ivanov, and S. Lajara. On moduli of smoothness and squareness. *Journal of Convex Analysis*, 17:441–449, 2010.
- [12] M. I. Kadets and B. S. Mityagin. Complemented subspaces in Banach spaces. *Russian Mathematical Surveys*(1973),28(6):77.
- [13] M.I. Kadets and M.G. Snobar. Some functionals over a compact Minkovskii space. *Mathematical notes of the Academy of Sciences of the USSR*, 10(4):694–696, 1971.
- [14] Peter M. Gruber. *Convex and Discrete Geometry*. Springer Berlin Heidelberg, 2007.
- [15] T. Figiel. On the moduli of convexity and smoothness. *Studia Math*, 56(2):121–155, 1976.
- [16] J. M. Borwein and J. D. Vanderwerff. *Convex Functions: Constructions, Characterizations and Counterexamples (Encyclopedia of Mathematics and its Applications)*. Cambridge University Press, 2010.
- [17] D. Amir. *Characterizations of Inner Product Spaces*. Birkhauser Verlag, 1986.
- [18] J. Diestel. *Geometry of Banach Spaces - Selected Topics*, volume 485. Springer-Verlag Berlin Heidelberg, 1975.
- [19] G. M. Ivanov. *Modulus of supporting convexity and supporting smoothness*. arXiv:1503.08912.
- [20] S. V. Konyagin. *On approximative properties of arbitrary closed sets in Banach spaces*. *Fund. Prikl. Mat.*, 3 (4) (1997), pp. 979–989 (in Russian)

DEPARTMENT OF HIGHER MATHEMATICS, MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, INSTITUTSKII PEREULOK 9, DOLGOPRUDNY, MOSCOW REGION, 141700, RUSSIA

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, SCHOOL OF APPLIED MATHEMATICS AND INFORMATION SCIENCE, BOLSHOI TREKHSVYATITELSKIY 3, MOSCOW, 109028, RUSSIA

E-mail address: grimivanov@gmail.com